# -CQG-IOP-

# Sample article for CQG-IOP-test2

Jane E  $Doe^{1*\dagger}$  and John RS Smith<sup>1,2</sup>

\*Correspondence: jane.e.doe@cambridge.co.uk <sup>1</sup>Department of Zoology, Cambridge, Waterloo Road, London, UK Full list of author information is available at the end of the article †Equal contributor

## Abstract

First part title: Text for this section.Second part title: Text for this section.Keywords: sample; article; author

## Content

Text and results for this section, as per the individual journal's instructions for authors.

## Section title

Text for this section ...

Sub-heading for section Text for this sub-heading ....

Sub-sub heading for section Text for this sub-sub-heading ...

Sub-sub-sub heading for section Text for this sub-sub-sub-heading ... In this section we examine the growth rate of the mean of  $Z_0$ ,  $Z_1$  and  $Z_2$ . In addition, we examine a common modeling assumption and note the importance of considering the tails of the extinction time  $T_x$  in studies of escape dynamics. We will first consider the expected resistant population at  $vT_x$  for some v > 0, (and temporarily assume  $\alpha = 0$ )

$$E[Z_1(vT_x)] = E\left[\mu T_x \int_0^{v \wedge 1} Z_0(uT_x) \exp(\lambda_1 T_x(v-u)) du\right]$$

If we assume that sensitive cells follow a deterministic decay  $Z_0(t) = xe^{\lambda_0 t}$  and approximate their extinction time as  $T_x \approx -\frac{1}{\lambda_0} \log x$ , then we can heuristically estimate the expected value as

$$E[Z_{1}(vT_{x})] = \frac{\mu}{r} \log x \int_{0}^{v \wedge 1} x^{1-u} x^{(\lambda_{1}/r)(v-u)} du$$
  
$$= \frac{\mu}{r} x^{1-\lambda_{1}/\lambda_{0}v} \log x \int_{0}^{v \wedge 1} x^{-u(1+\lambda_{1}/r)} du$$
  
$$= \frac{\mu}{\lambda_{1}-\lambda_{0}} x^{1+\lambda_{1}/rv} \left(1 - \exp\left[-(v \wedge 1)\left(1 + \frac{\lambda_{1}}{r}\right)\log x\right]\right).$$
(1)

Thus we observe that this expected value is finite for all v > 0 (also see [1, 2, 3, 4, 5]).

#### **Competing interests**

The authors declare that they have no competing interests.

#### Author's contributions

Text for this section ...

#### Acknowledgements

Text for this section ...

#### Author details

<sup>1</sup>Department of Zoology, Cambridge, Waterloo Road, London, UK. <sup>2</sup>Marine Ecology Department, Institute of Marine Sciences Kiel, Düsternbrooker Weg 20, 24105 Kiel, Germany.

#### References

- 1. Koonin, E.V., Altschul, S.F., Bork, P.: Brca1 protein products: functional motifs. Nat Genet 13, 266-267 (1996)
- 2. Kharitonov, S.A., Barnes, P.J.: Clinical Aspects of Exhaled Nitric Oxide. in press
- Zvaifler, N.J., Burger, J.A., Marinova-Mutafchieva, L., Taylor, P., Maini, R.N.: Mesenchymal cells, stromal derived factor-1 and rheumatoid arthritis [abstract]. Arthritis Rheum 42, 250 (1999)
- Jones, X.: Zeolites and synthetic mechanisms. In: Smith, Y. (ed.) Proceedings of the First National Conference on Porous Sieves: 27-30 June 1996; Baltimore, pp. 16–27 (1996). Stoneham: Butterworth-Heinemann
- 5. Margulis, L.: Origin of Eukaryotic Cells. Yale University Press, New Haven (1970)

#### Figures

Figure 1 Sample figure title. A short description of the figure content should go here.

Figure 2 Sample figure title. Figure legend text.

### Tables

 Table 1 Sample table title. This is where the description of the table should go.

	B1	B2	B3
A1	0.1	0.2	0.3
A2			
A3			

#### Additional Files

Additional file 1 — Sample additional file title Additional file descriptions text (including details of how to view the file, if it is in a non-standard format or the file extension). This might refer to a multi-page table or a figure.

Additional file 2 — Sample additional file title Additional file descriptions text.