# -DIR-

# Sample article for DIR

Jane E Doe<sup>1\*†</sup> and John RS Smith<sup>1,2</sup>

\*Correspondence:
jane.e.doe@cambridge.co.uk

¹Department of Zoology,
Cambridge, Waterloo Road,
London, UK

Full list of author information is
available at the end of the article
†Equal contributor

#### **Abstract**

**First part title:** Text for this section. **Second part title:** Text for this section.

Keywords: sample; article; author

## Content

Text and results for this section, as per the individual journal's instructions for authors.

## Section title

Text for this section ...

Sub-heading for section

Text for this sub-heading ...

Sub-sub heading for section

Text for this sub-sub-heading ...

Sub-sub-sub heading for section Text for this sub-sub-heading ... In this section we examine the growth rate of the mean of  $Z_0$ ,  $Z_1$  and  $Z_2$ . In addition, we examine a common modeling assumption and note the importance of considering the tails of the extinction time  $T_x$  in studies of escape dynamics. We will first consider the expected resistant population at  $vT_x$  for some v > 0, (and temporarily assume  $\alpha = 0$ )

$$E[Z_1(vT_x)] = E\left[\mu T_x \int_0^{v \wedge 1} Z_0(uT_x) \exp(\lambda_1 T_x(v-u)) du\right].$$

If we assume that sensitive cells follow a deterministic decay  $Z_0(t) = xe^{\lambda_0 t}$  and approximate their extinction time as  $T_x \approx -\frac{1}{\lambda_0} \log x$ , then we can heuristically estimate the expected value as

$$E[Z_1(vT_x)] = \frac{\mu}{r} \log x \int_0^{v \wedge 1} x^{1-u} x^{(\lambda_1/r)(v-u)} du$$

$$= \frac{\mu}{r} x^{1-\lambda_1/\lambda_0 v} \log x \int_0^{v \wedge 1} x^{-u(1+\lambda_1/r)} du$$

$$= \frac{\mu}{\lambda_1 - \lambda_0} x^{1+\lambda_1/r v} \left(1 - \exp\left[-(v \wedge 1)\left(1 + \frac{\lambda_1}{r}\right)\log x\right]\right). \quad (1)$$

Doe and Smith Page 2 of 2

Thus we observe that this expected value is finite for all v > 0 (also see [1, 2, 3, 4, 5]).

#### Competing interests

The authors declare that they have no competing interests.

#### Author's contributions

Text for this section ...

#### Acknowledgements

Text for this section ...

#### Author details

<sup>1</sup>Department of Zoology, Cambridge, Waterloo Road, London, UK. <sup>2</sup>Marine Ecology Department, Institute of Marine Sciences Kiel, Düsternbrooker Weg 20, 24105 Kiel, Germany.

#### References

- 1. Koonin, E.V., Altschul, S.F., Bork, P.: Brca1 protein products: functional motifs. Nat Genet 13, 266-267 (1996)
- 2. Kharitonov, S.A., Barnes, P.J.: Clinical Aspects of Exhaled Nitric Oxide. in press
- 3. Zvaifler, N.J., Burger, J.A., Marinova-Mutafchieva, L., Taylor, P., Maini, R.N.: Mesenchymal cells, stromal derived factor-1 and rheumatoid arthritis [abstract]. Arthritis Rheum 42, 250 (1999)
- 4. Jones, X.: Zeolites and synthetic mechanisms. In: Smith, Y. (ed.) Proceedings of the First National Conference on Porous Sieves: 27-30 June 1996; Baltimore, pp. 16–27 (1996). Stoneham: Butterworth-Heinemann
- 5. Margulis, L.: Origin of Eukaryotic Cells. Yale University Press, New Haven (1970)

#### **Figures**

Figure 1 Sample figure title. A short description of the figure content should go here.

Figure 2 Sample figure title. Figure legend text.

#### **Tables**

Table 1 Sample table title. This is where the description of the table should go.

	В1	B2	B3
A1	0.1	0.2	0.3
A2			
А3			

## **Additional Files**

Additional file 1 — Sample additional file title

Additional file descriptions text (including details of how to view the file, if it is in a non-standard format or the file extension). This might refer to a multi-page table or a figure.